

Image not found

Congratulations to New ICFO PhD Graduate

Dr. Laura Zarraoa Sardon graduated with a thesis entitled *Photon counting with a single neutral atom: quantum efficiency, dark counts, and background rejection*

February 27, 2025

We congratulate Dr. Laura Zarraoa Sardon who defended her thesis this morning in ICFO's Auditorium.

Dr. Zarraoa Sardon obtained her MSc in Physics and Nanotechnology from the Technical University of Denmark and she joined the Atomic Quantum Optics team led by ICREA Prof. Dr. Morgan Mitchell. Her thesis titled *Photon counting with a single neutral atom: quantum efficiency, dark counts, and background rejection* was supervised by ICREA Prof Dr. Morgan Mitchell and Dr. Romain Ve

-

ABSTRACT:

-

This thesis studies the use of a single trapped neutral ^{87}Rb atom as a photon counter. Detection of quantum jumps (QJs), i.e., abrupt changes between atomic states observable by a change in atomic fluorescence, is used to infer the arrival of single photons. This is referred to as the quantum jump photodetection (QJPD) technique.

The thesis first situates QJPD in the context of photodetection. Compared to traditional detectors, QJPD technique has lower speed and quantum efficiency (QE), but has exceptional performance in other figures of merit: QJPD is intrinsically narrowband, has strong out-of-band rejection, and very low dark counts (DCs). These features make the QJPD interesting for applications that detect weak optical signals in the presence of a strong broadband background.

Experimental methods to study QJPD are described. A ^{87}Rb atom is loaded from a magneto-optical trap (MOT) into a far-off resonance trap (FORT) at the center of four orthogonal, co-focal, high numerical aperture lenses. These lenses create the FORT, couple probe light onto the atom and collect the atomic fluorescence, which is used to identify the atomic state. A typical QJPD sequence is presented, which consists of trapping and cooling an atom in the FORT, optically pumping it into the dark state, illuminating with probe light, illuminating with readout light and collecting fluorescence photons, and checking that the

atom has not left the FORT during the sequence.

Statistical methods for measurement of QE and DC contributions are introduced. These compare the observed fluorescence count distribution against measured hyperfine-state fluorescence distributions. A QE of $(2.4 \pm 0.1) \times 10^{-3}$ is demonstrated, a record for single photon absorption by a single atom in free space.

Dark count contributions are measured. To produce low DC, the QJPD technique is implemented in two time windows: an exposure time for the single photon absorption followed by a short fluorescence time to read out the atomic state. This implies distinct acquisition and readout DC contributions, similarly to CCD and CMOS detectors. A dark jump rate (analogous to CCD/CMOS dark current) is measured of $(5 \pm 10) \times 10^{-3}$ jumps/s, consistent with zero and limited by measurement statistics. The measured readout contribution is $(4.0 \pm 0.4) \times 10^{-3}$ jumps per ms of fluorescence readout. For a 1 Hz readout rate, with 1 ms readout pulses, a net dark count rate of $(15 \pm 10) \times 10^{-3}$ counts/acquisition is demonstrated, which is already competitive with any non-cryogenic detector. The background rejection capabilities of the system are tested by measuring quantum jump rates when the atom is illuminated with direct sunlight, and with light scattered by the atmosphere (skylight). A rate equation model is developed to describe QJ probabilities in the presence of both intense broadband background and weak resonant probe light. This model is used to interpret experiments in which a weak signal beam competes with strong broadband background and validated using direct sunlight. Measurements where the atom is illuminated with skylight show no observable background-induced QJs. Finally, measurements of sky brightness and its fluctuations are presented, showing large fluctuations even on mostly clear days, a factor that further increases the need for background rejection. A number of contemporary applications of extreme photodetection, including free-space quantum communication in daylight, classical optical communications in space, and fundamental physics experiments, are discussed as possible applications of the QJPD technique. A realistic scenario where the demonstrated QJPD capabilities surpass the current performance of commercial single photodetectors is presented. Finally, potential improvements are discussed. It is shown that existing atomic and optical technologies could be applied to reach different wavelength ranges, narrower bandwidths, higher quantum efficiency, and lower dark

Thesis Committee:

Prof. Dr. Wolf von Klitzing, Institute of Electronic Structure and Laser

Prof. Dr. Hugues de Riedmatten, ICFO

Prof. Dr. Mathilde Hugbart, Institut de Physique de Nice